

Bruce T. Dr. Jet Propulsion Laboratory MS 169-506 CA 91109 Pasadena
USA
R.G. Marsden and E. Marsch

EGS Abstract for The Hague, 1999

MICROSTRUCTURES IN THE POLAR SOLAR WIND: ULYSSES

B. T. Tsurutani, J.K. Arballo, C. Galvan, B.E. Goldstein, G.S. Lakhina, R. Sakurai, E.J. Smith and M. Neugebauer
Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA.

bruce.tsurutani@jpl.nasa.gov/Fax: 818 354-8895

We find that small (10-200 rP) magnetic decreases comprise a dominant part of the polar solar wind microstructure at Ulysses distances (2.2 AU). These magnetic field dips are almost always bounded by tangential discontinuities, a feature which is not well understood at this time. Hundreds of these events have been examined in detail and a variety of types have been found. These will be described. It is speculated that these structures have been generated by perpendicular heating of ions closer to the Sun and have then been convected to distances of Ulysses. Such structures may be very important for the rapid cross- field diffusion of ions in the polar regions of the heliosphere.

Submittal Information

- | | |
|--|--|
| 1. Tsurutani
4800 Oak Grove Drive
Tel.: 818 354-7559
Fax.: [818 354-8895
E-mail: bruce.tsurutani@jpl.nasa.gov | 2. Session ST5 Open Session on Solar and
Heliospheric Physics
3.
4. none
5. Oral presentation strongly preferred |
|--|--|

Abstracts to be submitted on or before December 15, 1997 to

EGS Office
Max-Planck-Str. 13
37191 Katlenburg-Lindau
Germany

Tel.: [+49] 5556-1440
Fax.: [+49] 5556-4709
Email: **EGS@Copernicus.org**
<http://www.copernicus.org/EGS/EGS.html>